

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- ✓ TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS


IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

5

CLAIMS

1. A compound of the formula



or a pharmaceutically acceptable salt thereof, wherein:

10


R^1 is H, hydroxy or methoxy;

R^2 is hydroxy;

R^3 is C_1 - C_{10} alkyl, C_2 - C_{10} alkenyl, C_2 - C_{10} alkynyl, cyano, $-CH_2S(O)_nR^8$ wherein n is an integer ranging from 0 to 2, $-CH_2OR^8$, $-CH_2N(OR^8)R^8$, $-CH_2NR^8R^{15}$, $-(CH_2)_m(C_6-C_{10}$ aryl), or $-(CH_2)_m$ (5-10 membered heteroaryl), wherein m is an integer ranging from 0 to 4, and wherein the

15 foregoing R^3 groups are optionally substituted by 1 to 3 R^{16} groups;

or R^2 and R^3 are taken together to form an oxazoli ring as shown below



R^4 is H, $-C(O)R^9$, $-C(O)OR^9$, $-C(O)NR^9R^{10}$ or a hydroxy protecting group;

20 R^5 is $-SR^8$, $-(CH_2)_nC(O)R^8$ wherein n is 0 or 1, C_1 - C_{10} alkyl, C_2 - C_{10} alkenyl, C_2 - C_{10} alkynyl, $-(CH_2)_m(C_6-C_{10}$ aryl), or $-(CH_2)_m$ (5-10 membered heteroaryl), wherein m is an integer ranging from 0 to 4, and wherein the foregoing R^5 groups are optionally substituted by 1 to 3 R^{16} groups;

each R^6 and R^7 is independently H, hydroxy, C_1 - C_6 alkoxy, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, $-(CH_2)_m(C_6-C_{10}$ aryl), or $-(CH_2)_m$ (5-10 membered heteroaryl), wherein m is an integer ranging from 0 to 4;

- 5 each R^8 is independently H, C_1 - C_{10} alkyl, C_2 - C_{10} alkenyl, C_2 - C_{10} alkynyl,
 $-(CH_2)_q CR^{11} R^{12} (CH_2)_r NR^{13} R^{14}$ wherein q and r are each independently an integer ranging from 0
to 3 except q and r are not both 0, $-(CH_2)_m (C_6$ - C_{10} aryl), or $-(CH_2)_m$ (5-10 membered heteroaryl),
wherein m is an integer ranging from 0 to 4, and wherein the foregoing R^8 groups, except H, are
optionally substituted by 1 to 3 R^{16} groups;
- 10 or where R^8 is as $-CH_2 NR^9 R^{15}$, R^{15} and R^8 may be taken together to form a 4-10
membered monocyclic or polycyclic saturated ring or a 5-10 membered heteroaryl ring, wherein
said saturated and heteroaryl rings optionally include 1 or 2 heteroatoms selected from O, S and -
 $N(R^8)$ -, in addition to the nitrogen to which R^{15} and R^8 are attached, said saturated ring optionally
includes 1 or 2 carbon-carbon double or triple bonds, and said saturated and heteroaryl rings are
15 optionally substituted by 1 to 3 R^{16} groups;
- each R^9 and R^{10} is independently H or C_1 - C_6 alkyl;
- each R^{11} , R^{12} , R^{13} and R^{14} is independently selected from H, C_1 - C_{10} alkyl, $-(CH_2)_m (C_6$ - C_{10}
aryl), and $-(CH_2)_m$ (5-10 membered heteroaryl), wherein m is an integer ranging from 0 to 4, and
wherein the foregoing R^{11} , R^{12} , R^{13} and R^{14} groups, except H, are optionally substituted by 1 to 3
20 R^{16} groups;
- or R^{11} and R^{13} are taken together to form $-(CH_2)_p$ - wherein p is an integer ranging from 0
to 3 such that a 4-7 membered saturated ring is formed that optionally includes 1 or 2 carbon-
carbon double or triple bonds;
- or R^{13} and R^{14} are taken together to form a 4-10 membered monocyclic or polycyclic
25 saturated ring or a 5-10 membered heteroaryl ring, wherein said saturated and heteroaryl rings
optionally include 1 or 2 heteroatoms selected from O, S and $-N(R^8)$ -, in addition to the nitrogen to
which R^{13} and R^{14} are attached, said saturated ring optionally includes 1 or 2 carbon-carbon
double or triple bonds, and said saturated and heteroaryl rings are optionally substituted by 1 to 3
 R^{16} groups;
- 30 R^{15} is H, C_1 - C_{10} alkyl, C_2 - C_{10} alkenyl, or C_2 - C_{10} alkynyl, wherein the foregoing R^{15} groups
are optionally substituted by 1 to 3 substituents independently selected from halo and $-OR^9$;
- each R^{16} is independently selected from halo, cyano, nitro, trifluoromethyl, azido,
 $-C(O)R^{17}$, $-C(O)OR^{17}$, $-C(O)OR^{17}$, $-OC(O)OR^{17}$, $-NR^6C(O)R^7$, $-C(O)NR^6R^7$, $-NR^6R^7$, hydroxy, C_1 -
 C_6 alkyl, C_1 - C_6 alkoxy, $-(CH_2)_m (C_6$ - C_{10} aryl), and $-(CH_2)_m$ (5-10 membered heteroaryl), wherein m
35 is an integer ranging from 0 to 4, and wherein said aryl and heteroaryl substituents are optionally
substituted by 1 or 2 substituents independently selected from halo, cyano, nitro, trifluoromethyl,
azido, $-C(O)R^{17}$, $-C(O)OR^{17}$, $-C(O)OR^{17}$, $-OC(O)OR^{17}$, $-NR^6C(O)R^7$, $-C(O)NR^6R^7$, $-NR^6R^7$,
hydroxy, C_1 - C_6 alkyl, and C_1 - C_6 alkoxy;
- each R^{17} is independently selected from H, C_1 - C_{10} alkyl, C_2 - C_{10} alkenyl, C_2 - C_{10} alkynyl,

- 5 $-(CH_2)_m(C_6-C_{10} \text{ aryl})$, and $-(CH_2)_m(5-10 \text{ membered heteroaryl})$, wherein m is an integer ranging from 0 to 4;

with the proviso that R^6 is not H where R^3 is $-CH_2S(O)_nR^6$.

2. The compound of claim 1 wherein R^4 is H, acetyl, or benzyloxycarbonyl.

3. The compound of claim 2 wherein R^1 is hydroxy, R^2 is hydroxy, R^3 is $-CH_2NR^{15}R^6$ or
10 $-CH_2SR^6$.

4. The compound of claim 3 wherein R^3 is $-CH_2NR^{15}R^6$ and R^{15} and R^6 are independently selected from H, C_1-C_{10} alkyl, C_2-C_{10} alkenyl, and C_2-C_{10} alkynyl, wherein the foregoing R^{15} and R^6 groups, except H, are optionally substituted by 1 or 2 substituents independently selected from hydroxy, halo and C_1-C_6 alkoxy.

- 15 5. The compound of claim 4 wherein R^{15} and R^6 are each independently selected from H, methyl, ethyl, allyl, *n*-butyl, isobutyl, 2-methoxyethyl, cyclopentyl, 3-methoxypropyl, 3-ethoxypropyl, *n*-propyl, isopropyl, 2-hydroxyethyl, cyclopropyl, 2,2,2-trifluoroethyl, 2-propynyl, *sec*-butyl, *tert*-butyl, and *n*-hexyl.

6. The compound of claim 2 wherein R^1 is hydroxy, R^2 is hydroxy, R^3 is $-CH_2NHR^6$, and
20 R^6 is $-(CH_2)_m(C_6-C_{10} \text{ aryl})$ wherein m is an integer ranging from 0 to 4.

7. The compound of claim 6 wherein R^6 is phenyl or benzyl.

8. The compound of claim 2 wherein R^1 is hydroxy, R^2 is hydroxy, R^3 is $-CH_2NR^{15}R^6$, and R^{15} and R^6 are taken together to form a 4-10 membered saturated ring.

9. The compound of claim 8 wherein R^{15} and R^6 are taken together to form a piperidino,
25 trimethyleneimino, or morpholino ring.

10. The compound of claim 2 wherein R^1 is hydroxy, R^2 is hydroxy, R^3 is $-CH_2NR^{15}R^6$, and R^{15} and R^6 are taken together to form a 5-10 membered heteroaryl ring optionally substituted by 1 or 2 C_1-C_6 alkyl groups.

11. The compound of claim 10 wherein R^{15} and R^6 are taken together to form a
30 pyrrolidino, triazolyl, or imidazolyl ring wherein said heteroaryl groups are optionally substituted by 1 or 2 methyl groups.

12. The compound of claim 2 wherein R^1 is hydroxy, R^2 is hydroxy, R^3 is $-CH_2SR^6$, and R^6 is selected from C_1-C_{10} alkyl, C_2-C_{10} alkenyl, and C_2-C_{10} alkynyl, wherein said R^6 groups are optionally substituted by 1 or 2 substituents independently selected from hydroxy, halo and C_1-C_6
35 alkoxy.

13. The compound of claim 12 wherein R^6 is methyl, ethyl, or 2-hydroxyethyl.

14. The compound of claim 2 wherein R^1 is hydroxy, R^2 is hydroxy, and R^3 is selected from C_1-C_{10} alkyl, C_2-C_{10} alkenyl, and C_2-C_{10} alkynyl, wherein said R^3 groups are optionally

- 5 substituted by 1 or 2 substituents independently selected from hydroxy, $-C(O)R^{17}$, $-NR^6R^7$, halo, cyano, azido, 5-10 membered heteroaryl, and C_1-C_6 alkoxy.

15 15. The compound of claim 14 wherein R^3 is methyl, allyl, vinyl, ethynyl, 1-methyl-1-propenyl, 3-methoxy-1-propynyl, 3-dimethylamino-1-propynyl, 2-pyridylethynyl, 1-propynyl, 3-hydroxy-1-propynyl, 3-hydroxy-1-propenyl, 3-hydroxypropyl, 3-methoxy-1-propenyl, 3-methoxypropyl, 1-propynyl, n-butyl, ethyl, propyl, 2-hydroxyethyl, azidomethyl, formylmethyl, 6-cyano-1-pentynyl, 3-dimethylamino-1-propenyl, or 3-dimethylaminopropyl.


16. The compound of claim 2 wherein R^1 is hydroxy, R^2 is hydroxy, and R^3 is $-(CH_2)_m(5-10 \text{ membered heteroaryl})$ wherein m is an integer ranging from 0 to 4.

17. The compound of claim 16 wherein R^3 is 2-thienyl, 2-pyridyl, 1-methyl-2-imidazolyl, 2-furyl, or 1-methyl-2-pyrrolyl.


18. The compound of claim 2 wherein R^1 is hydroxy, R^2 is hydroxy, and R^3 is $-(CH_2)_m(C_6-C_{10} \text{ aryl})$ wherein m is an integer ranging from 0 to 4.

19. The compound of claim 18 wherein R^3 is phenyl.

20 20. The compound of claim 2 wherein R^2 and R^3 are taken together to form an oxazolyl ring as shown below



21. The compound of claim 2 wherein R^3 is selected from the following:




25 wherein X^3 is O, S or $-N(R^{15})-$, R^9 and R^{15} are as defined in claim 1, and the $-OR^9$ group may be attached at any available carbon on the phenyl group.

22. A pharmaceutical composition for the treatment of a bacterial infection or a protozoa infection in a mammal, fish, or bird which comprises a therapeutically effective amount of a compound of claim 1 and a pharmaceutically acceptable carrier.

30 23. A method of treating a bacterial infection or a protozoa infection in a mammal, fish, or bird which comprises administering to said mammal, fish or bird a therapeutically effective amount of a compound of claim 1.

24. A method of preparing a compound of the formula



5


or a pharmaceutically acceptable salt thereof, wherein:

R^1 is H, hydroxy or methoxy;

R^2 is hydroxy;

R^3 is C_1 - C_{10} alkyl, C_2 - C_{10} alkenyl, C_2 - C_{10} alkynyl, cyano, $-CH_2S(O)_nR^8$ wherein n is an integer ranging from 0 to 2, $-CH_2OR^8$, $-CH_2N(OR^8)R^8$, $-CH_2NR^8R^{15}$, $-(CH_2)_m(C_6-C_{10} \text{ aryl})$, or $-(CH_2)_m(5-10 \text{ membered heteroaryl})$, wherein m is an integer ranging from 0 to 4, and wherein the foregoing R^3 groups are optionally substituted by 1 to 3 R^{16} groups;

or R^2 and R^3 are taken together to form an oxazoly ring as shown below



15

R^4 is H, $-C(O)R^9$, $-C(O)OR^9$, $-C(O)NR^9R^{10}$ or a hydroxy protecting group;

R^5 is $-SR^8$, $-(CH_2)_nC(O)R^8$ wherein n is 0 or 1, C_1 - C_{10} alkyl, C_2 - C_{10} alkenyl, C_2 - C_{10} alkynyl, $-(CH_2)_m(C_6-C_{10} \text{ aryl})$, or $-(CH_2)_m(5-10 \text{ membered heteroaryl})$, wherein m is an integer ranging from 0 to 4, and wherein the foregoing R^5 groups are optionally substituted by 1 to 3 R^{16} groups;

each R^6 and R^7 is independently H, hydroxy, C_1 - C_6 alkoxy, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, $-(CH_2)_m(C_6-C_{10} \text{ aryl})$, or $-(CH_2)_m(5-10 \text{ membered heteroaryl})$, wherein m is an integer ranging from 0 to 4;

each R^8 is independently H, C_1 - C_{10} alkyl, C_2 - C_{10} alkenyl, C_2 - C_{10} alkynyl, $-(CH_2)_qCR^{11}R^{12}(CH_2)_rNR^{13}R^{14}$ wherein q and r are each independently an integer ranging from 0 to 3 except q and r are not both 0, $-(CH_2)_m(C_6-C_{10} \text{ aryl})$, or $-(CH_2)_m(5-10 \text{ membered heteroaryl})$,

5 wherein m is an integer ranging from 0 to 4, and wherein the foregoing R^8 groups, except H, are optionally substituted by 1 to 3 R^{16} groups;

or where R^8 is as $-CH_2NR^8R^{15}$, R^{15} and R^8 may be taken together to form a 4-10 membered monocyclic or polycyclic saturated ring or a 5-10 membered heteroaryl ring, wherein said saturated and heteroaryl rings optionally include 1 or 2 heteroatoms selected from O, S and -

10 $N(R^8)-$, in addition to the nitrogen to which R^{15} and R^8 are attached, said saturated ring optionally includes 1 or 2 carbon-carbon double or triple bonds, and said saturated and heteroaryl rings are optionally substituted by 1 to 3 R^{16} groups;

each R^9 and R^{10} is independently H or C_1-C_6 alkyl;

each R^{11} , R^{12} , R^{13} and R^{14} is independently selected from H, C_1-C_{10} alkyl, $-(CH_2)_m(C_6-C_{10}$ aryl), and $-(CH_2)_m(5-10$ membered heteroaryl), wherein m is an integer ranging from 0 to 4, and wherein the foregoing R^{11} , R^{12} , R^{13} and R^{14} groups, except H, are optionally substituted by 1 to 3 R^{16} groups;

or R^{11} and R^{13} are taken together to form $-(CH_2)_p-$ wherein p is an integer ranging from 0 to 3 such that a 4-7 membered saturated ring is formed that optionally includes 1 or 2 carbon-carbon double or triple bonds;

or R^{13} and R^{14} are taken together to form a 4-10 membered monocyclic or polycyclic saturated ring or a 5-10 membered heteroaryl ring, wherein said saturated and heteroaryl rings optionally include 1 or 2 heteroatoms selected from O, S and $-N(R^8)-$, in addition to the nitrogen to which R^{13} and R^{14} are attached, said saturated ring optionally includes 1 or 2 carbon-carbon double or triple bonds, and said saturated and heteroaryl rings are optionally substituted by 1 to 3 R^{16} groups;


R^{15} is H, C_1-C_{10} alkyl, C_2-C_{10} alkenyl, or C_2-C_{10} alkynyl, wherein the foregoing R^{15} groups are optionally substituted by 1 to 3 substituents independently selected from halo and $-OR^9$;

each R^{16} is independently selected from halo, cyano, nitro, trifluoromethyl, azido, $-C(O)R^{17}$, $-C(O)OR^{17}$, $-C(O)OR^{17}$, $-OC(O)OR^{17}$, $-NR^8C(O)R^7$, $-C(O)NR^8R^7$, $-NR^8R^7$, hydroxy, C_1-C_6 alkyl, C_1-C_6 alkoxy, $-(CH_2)_m(C_6-C_{10}$ aryl), and $-(CH_2)_m(5-10$ membered heteroaryl), wherein m is an integer ranging from 0 to 4, and wherein said aryl and heteroaryl substituents are optionally substituted by 1 or 2 substituents independently selected from halo, cyano, nitro, trifluoromethyl, azido, $-C(O)R^{17}$, $-C(O)OR^{17}$, $-C(O)OR^{17}$, $-OC(O)OR^{17}$, $-NR^8C(O)R^7$, $-C(O)NR^8R^7$, $-NR^8R^7$, hydroxy, C_1-C_6 alkyl, and C_1-C_6 alkoxy;

each R^{17} is independently selected from H, C_1-C_{10} alkyl, C_2-C_{10} alkenyl, C_2-C_{10} alkynyl, $-(CH_2)_m(C_6-C_{10}$ aryl), and $-(CH_2)_m(5-10$ membered heteroaryl), wherein m is an integer ranging from 0 to 4;

with the proviso that R^8 is not H where R^3 is $-CH_2S(O)_mR^8$;


- 5 which comprises treating a compound of the formula



wherein R^1 and R^4 are as defined above, with a compound of the formula HOR^8 , HSR^8 or $HNR^{15}R^8$, wherein n , R^{15} and R^8 are as defined above, wherein if said compound of formula HSR^8 is used the resulting R^2 group of formula $-CH_2SR^8$ is optionally oxidised to $-CH_2S(O)R^8$ or

- 10 $-CH_2S(O)_2R^8$.


25. The method of claim 24 wherein the compound of formula 5 is prepared by treating a compound of the formula



- wherein R^1 and R^4 are as defined in claim 24, with $(CH_3)_3S(O)_nX^2$, wherein n is 0 or 1 and
 15 X^2 is halo, $-BF_4$ or $-PF_6$, in the presence of a base.

26. The method of claim 25 wherein X^2 is iodo or BF_4 , and said base is selected from potassium tert-butoxide, sodium tert-butoxide, sodium ethoxide, sodium hydride, 1,1,3,3-tetramethylguanidine, 1,8-diazabicyclo[5.4.0]undec-7-ene, 1,5-diazabicyclo[4.3.0]non-5-ene, potassium hexamethyldisilazide (KHMDs), potassium ethoxide, and sodium methoxide.

27. A compound of the formula




or a pharmaceutically acceptable salt thereof, wherein:

R¹ is H, hydroxy or methoxy; and,

R⁴ is H, -C(O)R⁹, -C(O)OR⁹, -C(O)NR⁹R¹⁰ or a hydroxy protecting group; and, each R⁹ and R¹⁰ is independently H or C₁-C₆ alkyl.

28. A compound of the formula



5 or a pharmaceutically acceptable salt thereof, wherein:

R^1 is H, hydroxy or methoxy; and,

R^4 is H, $-C(O)R^9$, $-C(O)OR^9$, $-C(O)NR^9R^{10}$ or a hydroxy protecting group; and,

each R^9 and R^{10} is independently H or C_1 - C_6 alkyl.